

ARITHMETICAL RANK OF SQUAREFREE MONOMIAL IDEALS GENERATED BY FIVE ELEMENTS OR WITH ARITHMETIC DEGREE FOUR

KYOUKO KIMURA, GIANCARLO RINALDO, AND NAOKI TERAJ

ABSTRACT. Let I be a squarefree monomial ideal of a polynomial ring S . In this paper, we prove that the arithmetical rank of I is equal to the projective dimension of S/I when one of the following conditions is satisfied: (1) $\mu(I) \leq 5$; (2) $\text{arithdeg } I \leq 4$.

1. INTRODUCTION

Let S be a polynomial ring over a field K and I a squarefree monomial ideal of S . We denote by $G(I)$, the minimal set of monomial generators of I . The *arithmetical rank* of I , denoted by $\text{ara } I$, is defined by the minimal number r of elements $a_1, \dots, a_r \in S$ such that

$$\sqrt{(a_1, \dots, a_r)} = \sqrt{I}.$$

When the above equality holds, we say that a_1, \dots, a_r *generate I up to radical*. By definition, $\text{ara } I \leq \mu(I)$ holds, where $\mu(I)$ denotes the cardinality of $G(I)$. On the other hand, Lyubeznik [17] proved that

$$(1.1) \quad \text{pd}_S S/I \leq \text{ara } I,$$

where $\text{pd}_S S/I$ denotes the projective dimension of S/I over S . Since $\text{height } I \leq \text{pd}_S S/I$ always holds, we have

$$\text{height } I \leq \text{pd}_S S/I \leq \text{ara } I \leq \mu(I).$$

Then it is natural to ask when $\text{ara } I = \text{pd}_S S/I$ holds. Many authors investigated this problem; see [1, 2, 3, 4, 5, 6, 12, 14, 15, 16, 19, 20]. In particular in [14, 15], it was proved that the equality $\text{ara } I = \text{pd}_S S/I$ holds for the two cases that $\mu(I) - \text{height } I \leq 2$ and $\text{arithdeg } I - \text{indeg } I \leq 1$. Here $\text{arithdeg } I$ denotes the *arithmetical degree* of I , which is equal to the number of minimal primes of I , and $\text{indeg } I$ denotes the *initial degree* of I ; see Section 2. As a result we know that $\text{ara } I = \text{pd}_S S/I$ for squarefree monomial ideals I with $\mu(I) \leq 4$ or with $\text{arithdeg } I \leq 3$.

In this paper, hence, we concentrate our attention on the following two cases: $\mu(I) = 5$ and $\text{arithdeg } I = 4$. And the main result of this paper is as follows:

Theorem 1.1. *Let I be a squarefree monomial ideal of S . Suppose that I satisfies one of the following conditions:*

- (1) $\mu(I) \leq 5$.
- (2) $\text{arithdeg } I \leq 4$.

2000 *Mathematics Subject Classification.* 13F55.

Key words and phrases. monomial ideal, arithmetical rank, projective dimension.

Then

$$\text{ara } I = \text{pd}_S S/I.$$

Note that there exists an ideal I with $\mu(I) = 6$ such that $\text{ara } I > \text{pd}_S S/I$ when $\text{char } K \neq 2$; see [15, Section 6].

After we recall some definitions and properties of Stanley–Reisner ideals and hypergraphs in Sections 2 and 3, we give a combinatorial characterization for a squarefree monomial ideal I with $\text{pd}_S S/I = \mu(I) - 1$ using hypergraphs in Section 4. It is necessary because we must use the fact that projective dimension of S/I is characteristic-free for a squarefree monomial ideal I with $\mu(I) = 5$ in Section 6.

In the case that $\mu(I) \leq 4$ all the squarefree monomial ideals are classified using hypergraphs in [14, 15]. But in the case that $\mu(I) = 5$, $\text{height } I = 2$ and $\text{pd } S/I = 3$, which is an essential difficult part for our problem, a similar classification is practically impossible because of their huge number. According to a computer, there are about $2.3 \cdot 10^6$ hypergraphs corresponding to such ideals. We need a reduction. Hence we focus on the set of the most “general” members among them. We call it a *generic set*. In Section 5 we give a formal definition of a generic set and prove that it is enough to show $\text{ara } I = \text{pd}_S S/I$ for each member I of the generic set. But we cannot obtain the generic set in our case without a computer. In Section 6, we give an algorithm to find a generic set for the “connected” squarefree monomial ideals I with $\mu(I) = 5$, $\text{height } I = 2$ and $\text{pd } S/I = 3$ using *CoCoA* and *Nauty*. As a result of computation, we found that the generic set consists of just three ideals. In Section 7, we prove $\text{ara } I = \text{pd}_S S/I$ when $\mu(I) = 5$ by showing the same equality for all three members of the generic set.

Finally in Section 8, we focus on the squarefree monomial ideals I with $\text{arithdeg } I \leq 4$. Here we use another reduction. In terms of simplicial complexes as shown in [5, 12], we may remove a face with a free vertex from a simplicial complex Δ when we consider the problem whether $\text{ara } I_\Delta = \text{pd}_S S/I_\Delta$ holds. We translate it in terms of hypergraphs and after such reduction we show $\text{ara } I = \text{pd}_S S/I$ for remaining ideals I with $\text{arithdeg } I \leq 4$.

2. PRELIMINARIES

In this section, we recall some definitions and properties which are needed to prove Theorem 1.1.

Let M be a Noetherian graded S -module and

$$F_\bullet : 0 \longrightarrow \bigoplus_{j \geq 0} S(-j)^{\beta_{p,j}} \longrightarrow \cdots \longrightarrow \bigoplus_{j \geq 0} S(-j)^{\beta_{0,j}} \longrightarrow M \longrightarrow 0$$

a graded minimal free resolution of M over S , where $S(-j)$ is a graded free S -module whose k th piece is given by S_{k-j} . Then $\beta_{i,j} = \beta_{i,j}(M)$ is called a *graded Betti number* of M and $\beta_i = \sum_j \beta_{i,j}$ is called the i th (*total*) *Betti number* of M . The projective dimension of M over S is defined by p and denoted by $\text{pd}_S S/I$ or by $\text{pd } S/I$. The *initial degree* of M and the *regularity* of M are defined by

$$\text{indeg } M = \min\{j : \beta_{0,j}(M) \neq 0\}, \quad \text{reg } M = \max\{j - i : \beta_{i,j}(M) \neq 0\},$$

respectively.

Next, we recall some definitions and properties of Stanley–Reisner ideals, especially Alexander duality.

Let $V = [n] := \{1, 2, \dots, n\}$. A *simplicial complex* Δ on the vertex set V is a collection of subsets of V with the conditions (a) $\{v\} \in \Delta$ for all $v \in V$; (b) $F \in \Delta$ and $G \subset F$ imply $G \in \Delta$. An element of V is called a *vertex* of Δ and an element of Δ is called a *face*. A maximal face of Δ is called a *facet* of Δ . Let F be a face of Δ . The *dimension* of F , denoted by $\dim F$, is defined by $|F| - 1$, where $|F|$ denotes the cardinality of F . If $\dim F = i$, then F is called an i -face. The *dimension* of Δ is defined by $\dim \Delta := \max\{\dim F : F \in \Delta\}$. A simplicial complex which consists of all subsets of its vertex set is called a *simplex*. Let u be a new vertex and $F \subset V$. The *cone from u over F* is a simplex on the vertex set $F \cup \{u\}$; see [5, Definition 1, p. 3687]. We denote it by $\text{cone}_u F$. Then the union $\Delta \cup \text{cone}_u F$ is a simplicial complex on the vertex set $V \cup \{u\}$.

The *Alexander dual complex* Δ^* of Δ is defined by $\Delta^* = \{F \subset V : V \setminus F \notin \Delta\}$. If $\dim \Delta < n - 2$, then Δ^* is also a simplicial complex on the same vertex set V .

For a simplicial complex Δ on the vertex set $V = [n]$, we can associate a squarefree monomial ideal I_Δ of $S = K[x_1, \dots, x_n]$ which is generated by all products $x_{i_1} \cdots x_{i_s}$, $1 \leq i_1 < \cdots < i_s \leq n$ with $\{i_1, \dots, i_s\} \notin \Delta$. The ideal I_Δ is called the *Stanley–Reisner ideal* of Δ . It is well known that the minimal prime decomposition of I_Δ is given by

$$I_\Delta = \bigcap_{F \in \Delta: \text{facet}} P_F,$$

where $P_G = (x_i : i \in V \setminus G)$ for $G \subset V$. On the other hand, for a squarefree monomial ideal $I \subset S$ with $\text{indeg } I \geq 2$, there exists a simplicial complex Δ on the vertex set $V = [n]$ such that $I = I_\Delta$. If $\dim \Delta < n - 2$, i.e., $\text{height } I \geq 2$, then we can consider the squarefree monomial ideal $I^* := I_{\Delta^*}$, which is called the *Alexander dual ideal* of $I = I_\Delta$. Then

$$I^* = I_{\Delta^*} = (x^{V \setminus F} : F \in \Delta \text{ is a facet}),$$

where $x^G = \prod_{i \in G} x_i$ for $G \subset V$. It is easy to see that $I^{**} = I$, $\text{indeg } I^* = \text{height } I$, and $\text{arithdeg } I^* = \mu(I)$ hold. Moreover, the equality $\text{reg } I^* = \text{pd}_S S/I$ also holds; see [23, Corollary 1.6].

3. HYPERGRAPHS

For this section, we refer to Kimura, Terai and Yoshida [14], [15] for more detailed information.

Let $V = [\mu]$. A *hypergraph* \mathcal{H} on the vertex set V is a collection of subsets of V with $\bigcup_{F \in \mathcal{H}} F = V$. The definitions and notations of the vertex, face, and dimension are the same as those for a simplicial complex. We set $B(\mathcal{H}) = \{v \in V : \{v\} \in \mathcal{H}\}$ and $W(\mathcal{H}) = V \setminus B(\mathcal{H})$. For a hypergraph \mathcal{H} on a vertex set V , we define the i -*subhypergraph* of \mathcal{H} by $\mathcal{H}^i = \{F \in \mathcal{H} : \dim F = i\}$. We sometimes identify $B(\mathcal{H})$ with \mathcal{H}^0 . For $U \subset V(\mathcal{H})$, we define the restriction of a hypergraph \mathcal{H} to U by $\mathcal{H}_U = \{F \in \mathcal{H} : F \subset U\}$.

A hypergraph \mathcal{H} on the vertex set V is called *disconnected* if there exist hypergraphs $\mathcal{H}_1, \mathcal{H}_2 \subsetneq \mathcal{H}$ on vertex sets $V_1, V_2 \subsetneq V$, respectively such that $\mathcal{H}_1 \cup \mathcal{H}_2 = \mathcal{H}$, $V_1 \cup V_2 = V$, and $V_1 \cap V_2 = \emptyset$. A hypergraph which is not disconnected is called *connected*.

Let I be a squarefree monomial ideal of $S = K[x_1, \dots, x_n]$ with $G(I) = \{m_1, \dots, m_\mu\}$. We associate a hypergraph $\mathcal{H}(I)$ on the vertex set $V = [\mu]$ with I by setting

$$\mathcal{H}(I) := \{\{j \in V : m_j \text{ is divisible by } x_i\} : i = 1, 2, \dots, n\}.$$

Definition 3.1. Let F be a face of $\mathcal{H}(I)$. Then there exists a variable x of S such that

$$(3.1) \quad F = \{j \in V : m_j \text{ is divisible by } x\}.$$

We call a variable x of S with condition (3.1) a *defining variable* of F (in $\mathcal{H}(I)$).

Conversely, we say that a *variable x of S defines a face F* of $\mathcal{H}(I)$ if x is a defining variable of F .

Note that the choice of a defining variable is not necessarily unique. Since the minimal generators of a concrete squarefree monomial ideal I do not have indices, we usually regard $\mathcal{H}(I)$ as a hypergraph with unlabeled vertices. If $\mathcal{H}(I)$ can be regarded as a subhypergraph of $\mathcal{H}(J)$ as unlabeled hypergraphs for two squarefree monomial ideals I and J , we write $\mathcal{H}(I) \subset \mathcal{H}(J)$ by abuse of language.

On the other hand, we can construct a squarefree monomial ideal from a given hypergraph \mathcal{H} on the vertex set $V = [\mu]$ if \mathcal{H} satisfies the following *separability condition*:

For any two vertices $i, j \in V$,

there exist faces $F, G \in \mathcal{H}$ such that $i \in F \setminus G$ and $j \in G \setminus F$.

The way of construction is: first, we assign a squarefree monomial A_F to each face $F \in \mathcal{H}$ such that A_F and A_G are coprime if $F \neq G$. Then we set

$$I = \left(\prod_{\substack{F \in \mathcal{H} \\ j \in F}} A_F : j = 1, 2, \dots, \mu \right),$$

which is a squarefree monomial ideal with $\mathcal{H}(I) = \mathcal{H}$ by virtue of the separability. When we assign a variable x_F for each $F \in \mathcal{H}$, we write the corresponding ideal as $I_{\mathcal{H}}$ in $K[x_F : F \in \mathcal{H}]$.

For later use we prove the following proposition:

Proposition 3.2. *Let I, I' be squarefree monomial ideals of polynomial rings S, S' , respectively. Suppose that $\mu(I) = \mu(I')$ and $\mathcal{H}(I) \subset \mathcal{H}(I')$. Then we have $\text{ara } I \leq \text{ara } I'$.*

Proof. Set $\mathcal{H} = \mathcal{H}(I)$, $\mathcal{H}' = \mathcal{H}(I')$. We may assume that $I = I_{\mathcal{H}}$ and $I' = I_{\mathcal{H}'}$ with $\mathcal{H} \subset \mathcal{H}'$. Set $G(I') = \{m'_1, \dots, m'_\mu\}$. Let m_i for $i = 1, 2, \dots, \mu$ be the monomial obtained by substitution of 1 to x_F for $F \in \mathcal{H}' \setminus \mathcal{H}$ in m'_i . We may assume that $G(I) = \{m_1, \dots, m_\mu\}$. Assume that q'_1, \dots, q'_r generate I' up to radical. Let q_i for $i = 1, 2, \dots, r$ be the polynomial obtained by substitution of 1 to x_F for $F \in \mathcal{H}' \setminus \mathcal{H}$ in q'_i . We show that q_1, \dots, q_r generate I up to radical. Since $q'_i \in I'$ for $i = 1, 2, \dots, r$, we have $q_i \in I$. On the other hand, suppose $m_i^p \in (q'_1, \dots, q'_r)$ for some $p \geq 1$. Then we have $m_i^p \in (q_1, \dots, q_r)$. Hence q_1, \dots, q_r generate I up to radical. \square

4. SQUAREFREE MONOMIAL IDEALS WHOSE PROJECTIVE DIMENSION IS CLOSE TO THE NUMBER OF GENERATORS

Let $S = K[x_1, x_2, \dots, x_n]$ be the polynomial ring in n variables over a field K . We fix a squarefree monomial ideal $I = (m_1, m_2, \dots, m_\mu)$, where $G(I) = \{m_1, m_2, \dots, m_\mu\}$ is the minimal generating set of monomials for I . For a squarefree monomial ideal I , by the Taylor resolution of S/I we have $\text{pd}_S S/I \leq \mu(I)$. In this section we give a combinatorial characterization for the squarefree monomial ideal I with $\text{pd}_S S/I = \mu(I) - 1$ using hypergraphs.

First we consider the condition $\text{pd}_S S/I = \mu(I)$. Then the following proposition is easy and well known.

Proposition 4.1. *The following conditions are equivalent for a squarefree monomial ideal I :*

- (1) $\text{pd}_S S/I = \mu(I)$.
- (2) For the hypergraph $\mathcal{H} := \mathcal{H}(I)$ we have $B(\mathcal{H}) = V(\mathcal{H})$.

By Lyubeznik[17] we have $\text{pd}_S S/I = \text{cd}I := \max\{i : H_I^i(S) \neq 0\}$. Assuming that $\text{pd}_S S/I \leq \mu(I) - 1$, we have $\text{pd}_S S/I = \mu(I) - 1$ if and only if $H_I^{\mu-1}(S) \neq 0$. We give a combinatorial interpretation for the condition $H_I^{\mu-1}(S) \neq 0$.

Consider the following Čech complex:

$$\begin{aligned} C^\bullet &= \bigotimes_{i=1}^{\mu} (0 \longrightarrow S \longrightarrow S_{m_i} \longrightarrow 0) \\ &= 0 \longrightarrow S \xrightarrow{\delta^1} \bigoplus_{1 \leq i \leq \mu} S_{m_i} \xrightarrow{\delta^2} \bigoplus_{1 \leq i < j \leq \mu} S_{m_i m_j} \xrightarrow{\delta^3} \cdots \xrightarrow{\delta^\mu} S_{m_1 m_2 \cdots m_\mu} \longrightarrow 0. \end{aligned}$$

We describe δ^{r+1} as follows. Put $R := S_{m_{i_1} m_{i_2} \cdots m_{i_r}}$ and $\{j_1, j_2, \dots, j_s\} = \{1, 2, \dots, \mu\} \setminus \{i_1, i_2, \dots, i_r\}$, where $j_1 < j_2 < \cdots < j_s$ and $r + s = \mu$. Let $\psi_{j_p} : R \longrightarrow R_{m_{j_p}}$ be a natural map. For $u \in R$, we have

$$\delta^{r+1}(u) = \sum_{p=1}^s (-1)^{|\{q: i_q < j_p\}|} \psi_{j_p}(u) = \sum_{p=1}^s (-1)^{j_p - p} \psi_{j_p}(u).$$

For $F \subset [n]$, we define $x^F := \prod_{i \in F} x_i$. We define a simplicial complex $\Delta(F)$ by

$$\Delta(F) = \left\{ \{i_1, i_2, \dots, i_r\} \subset [\mu] : x^F \mid \prod_{j \in \{1, 2, \dots, \mu\} \setminus \{i_1, i_2, \dots, i_r\}} m_j \right\}.$$

For $\alpha \in \mathbf{Z}^n$, there is a unique decomposition $\alpha = \alpha_+ - \alpha_-$ such that $\alpha_+, \alpha_- \in \mathbf{N}^n$ and $\text{supp } \alpha_+ \cap \text{supp } \alpha_- = \emptyset$. Then we have $\text{supp } \alpha_- = \{i : \alpha_i < 0\}$.

Lemma 4.2 ([21, Lemma 2.5]). *For $\alpha \in \mathbf{Z}^n$ give an orientation for $\Delta(\text{supp } \alpha_-)$ by $1 < 2 < \cdots < \mu$. Then we have the following isomorphism of complexes: $C_\alpha^\bullet \cong \tilde{C}_\bullet(\Delta(\text{supp } \alpha_-))$ such that $C_\alpha^r \cong \tilde{C}_{\mu-r-1}(\Delta(\text{supp } \alpha_-))$.*

Then using the above lemma we have the following theorem:

Theorem 4.3. *The following conditions are equivalent for a squarefree monomial ideal I :*

- (1) $\text{pd}_S S/I = \mu(I) - 1$.
- (2) The hypergraph $\mathcal{H} := \mathcal{H}(I)$ satisfies $B(\mathcal{H}) \neq V(\mathcal{H})$ and either one of the following conditions:
 - (i) The graph $(W(\mathcal{H}), \mathcal{H}_{W(\mathcal{H})}^1)$ contains a complete bipartite graph as a spanning subgraph.
 - (ii) There exists $i \in B(\mathcal{H})$ such that $\{i, j\} \in \mathcal{H}$ for all $j \in W(\mathcal{H})$.

Proof. We may assume $B(\mathcal{H}) \neq V(\mathcal{H})$. By the above lemma we have the following isomorphisms:

$$\begin{aligned} H_I^{\mu-1}(S)_\alpha &= H^{\mu-1}(C^\bullet)_\alpha \\ &\cong \tilde{H}_0(\Delta(\text{supp } \alpha_-); K) \\ &\cong \tilde{H}_0(\Delta(\text{supp } \alpha_-)^{(1)}; K), \end{aligned}$$

where $\Delta(\text{supp } \alpha_-)^{(1)} := \{F \in \Delta(\text{supp } \alpha_-) : \dim F \leq 1\}$ is the 1-skeleton of $\Delta(\text{supp } \alpha_-)$. Hence $H_T^{\mu-1}(S)_\alpha = 0$ if and only if $\Delta(\text{supp } \alpha_-)^{(1)}$ is connected.

We claim that $\Delta(\text{supp } \alpha_-)^{(1)}$ is connected for all $\alpha \in \mathbf{Z}^n$ if and only if the graph $(U, \binom{U}{2} \setminus \mathcal{H}_U^1)$ is connected for all $W(\mathcal{H}) \subset U \subset V(\mathcal{H})$, where $\binom{U}{2} := \{\{i, j\} \subset U : i \neq j\}$. Let U be the vertex set of $\Delta(\text{supp } \alpha_-)^{(1)}$ for $\alpha \in \mathbf{Z}^n$. Then $U \supset W(\mathcal{H})$ and $\binom{U}{2} \setminus \mathcal{H}_U^1 \subset \Delta(\text{supp } \alpha_-)^{(1)}$. Hence if $(U, \binom{U}{2} \setminus \mathcal{H}_U^1)$ is connected, then so is $\Delta(\text{supp } \alpha_-)^{(1)}$. On the other hand, fix U such that $W(\mathcal{H}) \subset U \subset V(\mathcal{H})$. Put $U = W(\mathcal{H}) \cup B'$, where $B' \subset B(\mathcal{H})$. By a suitable change of variables, we may assume that $B' = \{1, 2, \dots, p\}$. For $1 \leq j \leq p$, set $\{x_i : x_i | m_j, x_i \nmid m_\ell \text{ for } 1 \leq \ell \leq \mu \text{ with } \ell \neq j\} = \{x_{i_{j_1}}, \dots, x_{i_{j_{s_j}}}\}$. Take $\alpha \in \mathbf{Z}^n$ such that $\text{supp } \alpha_- = [n] \setminus \{i_{11}, \dots, i_{1s_1}, i_{21}, \dots, i_{2s_2}, \dots, i_{p1}, \dots, i_{ps_p}\}$. Then we have $\binom{U}{2} \setminus \mathcal{H}_U^1$ is equal to the set of all 2-faces in $\Delta(\text{supp } \alpha_-)$. Hence we have the claim.

The graph $(U, \binom{U}{2} \setminus \mathcal{H}_U^1)$ is connected for all U such that $W(\mathcal{H}) \subset U \subset V(\mathcal{H})$ if and only if the following conditions (I) and (II) are satisfied:

- (I) The graph $(W(\mathcal{H}), \binom{W(\mathcal{H})}{2} \setminus \mathcal{H}_{W(\mathcal{H})}^1)$ is connected.
- (II) For $i \in B(\mathcal{H})$ set $U_i = W(\mathcal{H}) \cup \{i\}$. The graph $(U_i, \binom{U_i}{2} \setminus \mathcal{H}_{U_i}^1)$ is connected for all $i \in B(\mathcal{H})$.

Hence the condition $\text{pd}_S S/I = \mu(I) - 1$ holds if and only if one of the following conditions (i)' or (ii)' is satisfied:

- (i)' The graph $(W(\mathcal{H}), \binom{W(\mathcal{H})}{2} \setminus \mathcal{H}_{W(\mathcal{H})}^1)$ is disconnected.
- (ii)' The graph $(U_i, \binom{U_i}{2} \setminus \mathcal{H}_{U_i}^1)$ is disconnected for some $i \in B(\mathcal{H})$.

The condition (i)' is equivalent to the condition (i). Under the assumption that the condition (i) (or equivalently (i)') does not hold the conditions (ii) and (ii)' are equivalent. Hence we are done. \square

Corollary 4.4. *The condition $\text{pd}_S S/I = \mu(I) - 1$ is independent of the base field K for a monomial ideal I .*

5. GENERIC SET

We cannot restrict the number of variables in general when we classify squarefree monomial ideals with a certain given property. In that case, it is convenient to consider finitely generated squarefree monomial ideals in a polynomial ring with infinite variables.

Let K be a field and $S_\infty = K[x_1, x_2, \dots]$ a polynomial ring with countably infinite variables over K . Let \mathcal{I} be the set of all finitely generated squarefree monomial ideals of S_∞ . For $I \in \mathcal{I}$, we denote by $X(I)$ the set of all variables which appear in one of the minimal monomial generators of I .

Definition 5.1. Let \mathcal{C} be some property on \mathcal{I} . That is, the subset

$$\mathcal{I}(\mathcal{C}) := \{I \in \mathcal{I} : I \text{ satisfies the property } \mathcal{C}\}$$

is uniquely determined. A subset $\mathcal{A} \subset \mathcal{I}$ is called a *generic set* on \mathcal{C} if the following two conditions are satisfied:

- (1) $\mathcal{A} \subset \mathcal{I}(\mathcal{C})$.
- (2) For any $J \in \mathcal{I}(\mathcal{C})$ with $G(J) = \{m_1, \dots, m_\mu\}$, there exist $I \in \mathcal{A}$ with $X(I) = \{x_1, \dots, x_n\}$ and $G(I) = \{m'_1, \dots, m'_\mu\}$ where $m'_i = x_{t_{i1}} \cdots x_{t_{ij_i}}$,

and (possibly trivial) squarefree monomials M_1, \dots, M_n on $X(J)$ that are pairwise coprime such that

$$M_{t_{i1}} \cdots M_{t_{ij_i}} = m_i, \quad i = 1, 2, \dots, \mu.$$

We say that a generic set \mathcal{A} on \mathcal{C} is *minimal* if \mathcal{A} is minimal among generic sets on \mathcal{C} with respect to inclusion, and we say \mathcal{A} on \mathcal{C} is *reduced* if

$$K[x_F : F \in \mathcal{H}(I)]/I_{\mathcal{H}(I)} \cong K[X(I)]/(I \cap K[X(I)])$$

for all $I \in \mathcal{A}$.

A minimal generic set has the following property:

Proposition 5.2. *Let \mathcal{A} be a minimal generic set on some property \mathcal{C} .*

- (1) *For $J \in \mathcal{I}(\mathcal{C})$, there exists $I \in \mathcal{A}$ such that $\mathcal{H}(J) \subset \mathcal{H}(I)$ and $\mu(J) = \mu(I)$.*
- (2) *If $\mathcal{H}(I') \subset \mathcal{H}(I)$ and $\mu(I') = \mu(I)$ for $I', I \in \mathcal{A}$, then $I' = I$.*

Proof. (1) By definition, there exists an ideal $I \in \mathcal{A}$ with the condition (2) of Definition 5.1. In particular, $\mu(I) = \mu(J) =: \mu$. We use the same notations as in Definition 5.1. Take a face $F \in \mathcal{H}(J)$ and let y_F be a defining variable of F in $\mathcal{H}(J)$. Take $i \in F$. Then y_F divides m_i . Since m_i can be written as the product $M_{t_{i1}} \cdots M_{t_{ij_i}}$, we may assume that $M_{t_{i1}}$ is divisible by y_F . Then for $1 \leq \ell \leq \mu$, the variable y_F divides m_ℓ if and only if the variable $x_{t_{i1}}$ divides m'_ℓ . This means that $x_{t_{i1}}$ defines F of $\mathcal{H}(I)$.

(2) First note that in the proof of (1), we also proved that $\mathcal{H}(J) \subset \mathcal{H}(I)$ holds when I and J have the connection as in Definition 5.1 (2). Therefore from the minimality of \mathcal{A} , it is enough to prove that if $\mathcal{H}(J) \subset \mathcal{H}(I)$ with $\mu(J) = \mu(I) = \mu$ for $I, J \in \mathcal{I}$, then I and J have that connection.

Let $G(J) = \{m_1, \dots, m_\mu\}$ and $G(I) = \{m'_1, \dots, m'_\mu\}$. For $F \in \mathcal{H}(J)$ (resp. $G \in \mathcal{H}(I)$), we denote by M_F (resp. M'_G), the product of all defining variables of F in $\mathcal{H}(J)$ (resp. G in $\mathcal{H}(I)$). Then

$$m_i = \prod_{\substack{F \in \mathcal{H}(J) \\ i \in F}} M_F, \quad m'_i = \prod_{\substack{F \in \mathcal{H}(J) \\ i \in F}} M'_F \prod_{\substack{G \in \mathcal{H}(I) \setminus \mathcal{H}(J) \\ i \in G}} M'_G.$$

Since $F \in \mathcal{H}(I)$, the product $M'_F \neq 1$. Let x_F be a variable which divides M'_F . Then the substitution M_F to x_F ; 1 to the variables dividing one of M'_F/x_F , M'_G yields the desired connection. \square

By virtue of Propositions 3.2 and 5.2 we have the following proposition:

Proposition 5.3. *Let \mathcal{A} be a generic set on some property \mathcal{C} . Suppose for all $I, J \in \mathcal{I}(\mathcal{C})$ we have*

$$\text{pd}_{K[X(I)]} K[X(I)]/(I \cap K[X(I)]) = \text{pd}_{K[X(J)]} K[X(J)]/(J \cap K[X(J)]).$$

We also assume that

$$\text{ara}(I \cap K[X(I)]) = \text{pd}_{K[X(I)]} K[X(I)]/(I \cap K[X(I)])$$

for all $I \in \mathcal{A}$. Then we have

$$\text{ara}(I \cap K[X(I)]) = \text{pd}_{K[X(I)]} K[X(I)]/(I \cap K[X(I)])$$

for all $I \in \mathcal{I}(\mathcal{C})$.

Let I be a squarefree monomial ideal of S . The next proposition guarantees that we may assume that $\mathcal{H}(I)$ is connected when we consider the problem on the arithmetical rank.

Proposition 5.4. *Let I_1, I_2 be squarefree monomial ideals of S . Suppose that $X(I_1) \cap X(I_2) = \emptyset$. Then we have*

$$\text{ara}(I_1 + I_2) \leq \text{ara } I_1 + \text{ara } I_2.$$

Moreover, if $\text{ara } I_i = \text{pd}_S S/I_i$ holds for $i = 1, 2$, then $\text{ara}(I_1 + I_2) = \text{pd}_S S/(I_1 + I_2)$ also holds.

Proof. When $f_1, \dots, f_{s_1} \in S$ generate I_1 up to radical and $g_1, \dots, g_{s_2} \in S$ generate I_2 up to radical, it is clear that these $s_1 + s_2$ elements generate $I_1 + I_2$ up to radical. Hence, we have the desired inequality.

Let F_\bullet, G_\bullet be minimal free resolutions of $S/I_1, S/I_2$, respectively. Then $F_\bullet \otimes G_\bullet$ provides a minimal free resolution of $S/(I_1 + I_2)$, and its length is equal to

$$\text{pd}_S S/I_1 + \text{pd}_S S/I_2 = \text{ara } I_1 + \text{ara } I_2 \geq \text{ara}(I_1 + I_2)$$

when $\text{ara } I_i = \text{pd}_S S/I_i$ holds for $i = 1, 2$. Since the inequality $\text{ara}(I_1 + I_2) \geq \text{pd}_S S/(I_1 + I_2)$ always holds by (1.1), we have the desired equality. \square

Let I be a squarefree monomial ideal with $G(I) = \{m_1, \dots, m_\mu\}$. We say that I is *connected* if for any distinct indices $i, j \in [\mu]$, there exists a sequence of indices $i = i_0, i_1, \dots, i_\ell = j$ such that $\gcd(m_{i_{k-1}}, m_{i_k}) \neq 1$ for all $k = 1, \dots, \ell$. In other words, the corresponding hypergraph $\mathcal{H}(I)$ is connected.

Remark 5.5. Proposition 5.4 guarantees that we may assume $\text{indeg } I \geq 2$. If $\text{indeg } I = 1$, then I can be written as $I_1 + (x)$, where $x \notin X(I_1)$. Then $\text{pd}_S S/I = \text{pd}_S S/I_1 + 1$ and $\text{ara } I \leq \text{ara } I_1 + 1$. Therefore if $\text{ara } I_1 = \text{pd}_S S/I_1$ holds, then $\text{ara } I = \text{pd}_S S/I$ also holds.

6. AN ALGORITHM TO FIND A MINIMAL REDUCED GENERIC SET

In this section we present an algorithm for computing a minimal reduced generic set on the property:

$\mathcal{C} : \mu(I) = 5, \text{pd}_{K[X(I)]} K[X(I)]/(I \cap K[X(I)]) = 3, \text{height } I = 2, I \text{ is connected.}$

The algorithm is composed mainly by three steps:

- Step 1) List up all the connected hypergraphs with five vertices which satisfy the separability condition;
- Step 2) Choose the hypergraphs \mathcal{H} with height $I_{\mathcal{H}} = 2$ and with $\text{pd } K[X(I_{\mathcal{H}})]/I_{\mathcal{H}} = 3$;
- Step 3) Find a minimal reduced generic set on \mathcal{C} .

Since we use a computer we must note that the projective dimension does not depend on the characteristic of K when $\mu(I) = 5$.

Lemma 6.1. *Let I be a squarefree monomial ideal of S with $\mu(I) = 5$. The projective dimension $\text{pd}_S S/I$ does not depend on the characteristic of K .*

Proof. We fix the base field K . If $\text{pd}_S S/I = 1$, then I is a principal ideal and the claim is clear.

Suppose $\text{pd}_S S/I = 2$. Then the height of I is either 1 or 2. We assume that height $I = 1$. Then the total Betti numbers do not change if we remove the prime

components of height one from I . Hence we may assume that $\text{height } I = 2$. Since $\text{pd}_S S/I = \text{height } I$, S/I is Cohen-Macaulay. Since Cohen-Macaulayness of the height two monomial ideals does not depend on the base field, we are done in this case.

If $\text{pd}_S S/I \geq 4$, then $\text{pd}_S S/I$ does not depend on the characteristic of K by Proposition 4.1 and Corollary 4.4.

Finally, suppose $\text{pd}_S S/I = 3$. By the above argument it is not possible to have $\text{pd}_S S/I \neq 3$ for another base field. Then we are done. \square

Now we explain the details of each step.

Step 1): To list up all the connected hypergraphs with five vertices, we consider a correspondence between a hypergraph and a bipartite graph.

First we give necessary definitions.

Let G be a graph on the vertex set V with the edge set $E(G)$. For a vertex $v \in V$, the set of neighbourhoods of v is defined by $N(v) := \{u \in V : \{u, v\} \in E(G)\}$. For a subset $V' \subset V$, we write $G_{V'}$ the induced subgraph of G on V' . For a vertex set V we consider a labeled partition $V = X \cup Y$. It means that X and Y are labeled. We call X the *first subset* (or the *indeterminate-part subset*) of V and Y the *second subset* (or the *generator-part subset*) of V . Note that a bipartite graph with a labeled partition $V = X \cup Y$ can be regarded as a directed bipartite graph on the partition $V = X \cup Y$ such that every edge $\{x, y\}$ has the direction from $x \in X$ to $y \in Y$.

Definition 6.2. Set $X = \{x_1, \dots, x_n\}$, $Y = \{y_1, \dots, y_\mu\}$. A bipartite graph G with the labeled partition $V = X \cup Y$ is said to be a *corresponding graph* if it satisfies the following three conditions:

- (1) The graph G is connected;
- (2) $N(y_i) \not\subset N(y_j)$ for all $i \neq j$;
- (3) $N(x_i) \neq N(x_j)$ for any $i \neq j$.

Two corresponding graphs G and G' with labeled partitions $V = X \cup Y$ and $V' = X' \cup Y'$, respectively, are isomorphic if there are bijections $f : X \rightarrow X'$ and $g : Y \rightarrow Y'$ such that $\{x, y\} \in E(G)$ for $x \in X$ and $y \in Y$ if and only if $\{f(x), g(y)\} \in E(G')$.

Now we describe the correspondence more concretely.

Proposition 6.3. We set $X = \{x_1, \dots, x_n\}$, $Y = \{y_1, \dots, y_\mu\}$ and $V = X \cup Y$. Let \mathcal{G} be the set of all isomorphism classes of corresponding graphs with the labeled partition $V = X \cup Y$ and \mathcal{S} be the set of all isomorphism classes of connected hypergraphs with the vertex set Y and with n faces which satisfy the separability condition.

Then the map $\phi : \mathcal{G} \rightarrow \mathcal{S}$ defined by sending $G \in \mathcal{G}$ to $\mathcal{H} = \{F_1, \dots, F_n\} \in \mathcal{S}$ where $F_i = \{y_j : \{x_i, y_j\} \in E(G)\}$ gives a one-to-one correspondence between these two sets.

Proof. Let G be a corresponding graph on the labeled partition $V = X \cup Y$. The connectivity of G corresponds to that of the hypergraph $\phi(G)$. The condition (2) of Definition 6.2 corresponds to the separability condition for the hypergraph $\phi(G)$. Moreover the condition (3) of Definition 6.2 guarantees that the hypergraph $\phi(G)$ has n faces. Hence the map ϕ is well defined. A hypergraph $\mathcal{H} = \{F_1, \dots, F_n\}$

on the vertex set $Y = \{y_1, \dots, y_\mu\}$ with the separability condition is associated to the bipartite graph on the labeled partition $X \cup Y$, where $X = \{x_1, \dots, x_n\}$ with $E(G) = \{\{x_i, y_j\} : y_j \in F_i\}$. This correspondence gives the inverse map of ϕ . \square

By virtue of Proposition 6.3, it is enough to enumerate all the corresponding graphs with labeled partition $V = X \cup Y$ such that $|Y| = 5$ and $|X| \leq 2^5 = 32$ to list up the corresponding hypergraphs.

To perform this computation we used an existent software called *Nauty* (see [8]), whose main purpose is to calculate a set of non-isomorphic graphs in an efficient way. We also customized it by a routine in C language to obtain graphs satisfying the conditions in Definition 6.2. See the site [13] for technical information and examples of computation. In this site we also provide the source under GPL license to generate the software applications.

As a result, in the case $\mu = 5$ the number of the corresponding graphs, or equivalently the number of the 5-vertex hypergraphs with the separability condition, is around $1.8 \cdot 10^7$. We denote the set of the above graphs by \mathcal{P}_1 .

Step 2): To each element $G \in \mathcal{P}_1$, we associate the squarefree monomial ideal $I = (m_1, \dots, m_5)$ in $S = K[x_1, \dots, x_{32}]$, where $m_i = \prod_{x_j \in N(y_i)} x_j$ for $i = 1, \dots, 5$.

Then we can use *CoCoA* ([9]) to determine whether height $I = 2$ and $\text{pd } S/I = 3$. Choose all graphs in \mathcal{P}_1 which correspond to an ideal satisfying the above properties. Set this subset as \mathcal{P}_2 .

By this computation we know that the cardinality of \mathcal{P}_2 is around $2.3 \cdot 10^6$.

Step 3): For the last step of our algorithm we need to find a minimal generic set on the property \mathcal{C} , or, equivalently, the corresponding set of bipartite graphs. For this purpose we define a partial order on bipartite graphs:

Definition 6.4. Let G and G' be bipartite graphs on the labeled partitions $X \cup Y$ and $X' \cup Y$, respectively. We introduce a partial order \preceq by setting $G' \preceq G$ if and only if there is a subset X'' of X such that $G_{X'' \cup Y} \cong G'$.

We want to list up all the maximal corresponding graphs in \mathcal{P}_2 with respect to the partial order \preceq . To reach this goal we partition the set \mathcal{P}_2 of the corresponding graphs

$$\mathcal{P}_2 = \bigcup \mathcal{G}_i$$

where $G \in \mathcal{G}_i$ if the first subset X has the cardinality i . Let ℓ be the maximum i such that $\mathcal{G}_i \neq \emptyset$. Obviously each $G \in \mathcal{G}_\ell$ is a maximal graph. Set $\mathcal{M}_\ell = \mathcal{G}_\ell$.

As the next step we list up the maximal graphs in $\mathcal{G}_{\ell-1}$. For each graph $G \in \mathcal{M}_\ell$ and for each vertex $x_i \in X$ let $G'_i = G_{V(G) \setminus \{x_i\}}$. We make the list $\{G'_i : G \in \mathcal{G}_\ell, x_i \in X\}$. If a graph in $\mathcal{G}_{\ell-1}$ is isomorphic to some G'_i then we remove it. At the end of this process we have only the maximal graphs in $\mathcal{G}_{\ell-1}$. We denote the set of the maximal graphs in $\mathcal{G}_{\ell-1}$ by $\mathcal{M}_{\ell-1}$.

As the next step we similarly list up the maximal graphs in $\mathcal{G}_{\ell-2}$. For each graph $G \in \mathcal{G}_{\ell-1}$ and for each vertex $x_i \in X$ let $G'_i = G_{V(G) \setminus \{x_i\}}$. We make the list $\{G'_i : G \in \mathcal{G}_{\ell-1}, x_i \in X\}$. If a graph in $\mathcal{G}_{\ell-2}$ is isomorphic to a graph in the list, then we remove it. At the end of this process we have only the maximal graphs in $\mathcal{G}_{\ell-2}$. We denote the set of the maximal graphs in $\mathcal{G}_{\ell-2}$ by $\mathcal{M}_{\ell-2}$.

We repeat a similar procedure for each $i \leq \ell - 3$ and define \mathcal{M}_i as above.

Set

$$\mathcal{P}_3 = \bigcup \mathcal{M}_i.$$

Then \mathcal{P}_3 is the set of maximal corresponding graphs. A corresponding set of square-free monomial ideals gives a minimal generic set on \mathcal{C} .

As the result performing the algorithm we have $|\mathcal{P}_3| = 3$. More concretely, we have the following theorem:

Theorem 6.5. *Let \mathcal{C} be the property on \mathcal{I} such that*

$$\mathcal{I}(\mathcal{C}) = \left\{ I \in \mathcal{I} : \begin{array}{l} \mu(I) = 5, \text{ pd}_{K[X(I)]} K[X(I)]/(I \cap K[X(I)]) = 3, \\ \text{height } I = 2, \text{ } I \text{ is connected} \end{array} \right\}.$$

Then $\{J_1, J_2, J_3\}$ is a minimal reduced generic set on \mathcal{C} , where we define $J_i = (m_1, m_2, m_3, m_4, m_5)$, $i = 1, 2, 3$, as follows:

For J_1 where $|X(J_1)| = 23$:

$$\begin{aligned} m_1 &= x_1 x_3 x_7 x_8 x_9 x_{10} x_{12} x_{14} x_{15} x_{18} x_{19} x_{20} x_{21} x_{22}, \\ m_2 &= x_2 x_4 x_5 x_8 x_9 x_{11} x_{13} x_{14} x_{17} x_{18} x_{19} x_{20} x_{21} x_{23}, \\ m_3 &= x_3 x_5 x_9 x_{11} x_{12} x_{15} x_{16} x_{17} x_{19} x_{20} x_{22} x_{23}, \\ m_4 &= x_4 x_6 x_{10} x_{11} x_{12} x_{13} x_{16} x_{18} x_{19} x_{21} x_{22} x_{23}, \\ m_5 &= x_6 x_7 x_{10} x_{13} x_{14} x_{15} x_{16} x_{17} x_{20} x_{21} x_{22} x_{23}. \end{aligned}$$

For J_2 where $|X(J_2)| = 24$:

$$\begin{aligned} m_1 &= x_1 x_4 x_7 x_9 x_{10} x_{12} x_{15} x_{16} x_{18} x_{19} x_{20} x_{22} x_{23} x_{24}, \\ m_2 &= x_2 x_5 x_8 x_9 x_{11} x_{14} x_{15} x_{17} x_{18} x_{19} x_{21} x_{22} x_{23} x_{24}, \\ m_3 &= x_3 x_6 x_7 x_8 x_{11} x_{12} x_{13} x_{16} x_{17} x_{19} x_{20} x_{21} x_{23} x_{24}, \\ m_4 &= x_4 x_5 x_{10} x_{11} x_{12} x_{13} x_{14} x_{15} x_{20} x_{21} x_{22} x_{23}, \\ m_5 &= x_6 x_{10} x_{13} x_{14} x_{16} x_{17} x_{18} x_{20} x_{21} x_{22} x_{24}. \end{aligned}$$

For J_3 where $|X(J_3)| = 25$:

$$\begin{aligned} m_1 &= x_1 x_5 x_6 x_7 x_{11} x_{12} x_{15} x_{18} x_{19} x_{20} x_{21} x_{22} x_{23} x_{24}, \\ m_2 &= x_2 x_5 x_8 x_9 x_{11} x_{13} x_{15} x_{16} x_{17} x_{18} x_{21} x_{22} x_{24} x_{25}, \\ m_3 &= x_3 x_6 x_8 x_{10} x_{12} x_{14} x_{15} x_{16} x_{17} x_{20} x_{21} x_{23} x_{24} x_{25}, \\ m_4 &= x_4 x_7 x_9 x_{10} x_{13} x_{14} x_{17} x_{18} x_{19} x_{20} x_{22} x_{23} x_{24} x_{25}, \\ m_5 &= x_{11} x_{12} x_{13} x_{14} x_{16} x_{19} x_{21} x_{22} x_{23} x_{25}. \end{aligned}$$

Remark 6.6. We have performed a similar algorithm for a minimal generic set on the property:

$\mathcal{C} : \mu(I) = 5, \text{ pd}_{K[X(I)]} K[X(I)]/(I \cap K[X(I)]) = 3, \text{ height } I = 3, I \text{ is connected.}$

As a result, we know that a minimal generic set consists of 9 ideals, which is coincident with the list chosen from all corresponding hypergraphs in [15], where a combinatorial argument is used without a computer.

7. THE CASE $\mu(I) \leq 5$

In this section, we compute the arithmetical rank of a squarefree monomial ideal I with $\mu(I) \leq 5$.

Theorem 7.1. *Let I be a squarefree monomial ideal of $S = K[x_1, \dots, x_n]$. Suppose that $\mu(I) \leq 5$. Then*

$$\text{ara } I = \text{pd}_S S/I.$$

Proof. If height $I = 1$, then I is of the form

$$I = (x_1, \dots, x_t)I'S,$$

where I' is a squarefree monomial ideal of $S' = K[x_{t+1}, \dots, x_n]$ with height $I' \geq 2$. Then the equality $\text{ara } I = \text{pd}_S S/I$ holds if the same equality holds for $I' \subset S'$. Hence we may assume that height $I \geq 2$. By [14, 15], we have already known $\text{ara } I = \text{pd}_S S/I$ when $\mu(I) - \text{height } I \leq 2$ or $\mu(I) - \text{pd}_S S/I \leq 1$. The remain cases are that $\mu(I) = 5$, height $I = 2$, and $\text{pd}_S S/I = 2, 3$. If $\text{pd}_S S/I = 2$, then $\text{pd}_S S/I = \text{height } I = 2$, and thus S/I is Cohen–Macaulay. The equality $\text{ara } I = \text{pd}_S S/I$ for such an ideal was proved in [12]. Therefore we only need to consider the case that $\mu(I) = 5$, height $I = 2$, and $\text{pd}_S S/I = 3$. By Proposition 5.3 and Theorem 6.5, it is enough to prove $\text{ara } I = \text{pd}_S S/I$ for the following three cases:

- (1) $I = J_1 \cap K[X]$ with $|X| = 23$;
- (2) $I = J_2 \cap K[X]$ with $|X| = 24$;
- (3) $I = J_3 \cap K[X]$ with $|X| = 25$.

Since we have $\text{ara } I \geq \text{pd}_S S/I = 3$ by (1.1), it is sufficient to show $\text{ara } I \leq 3$.

Case (1): We prove that the following 3 elements g_1, g_2, g_3 generate I up to radical:

$$g_1 = x_{16}f_1f_2, \quad g_2 = m_1f_1 + m_4, \quad g_3 = m_2f_2 + m_5,$$

where

$$f_1 = \gcd(m_1, m_3) + \gcd(m_4, m_5)m_2, \quad f_2 = \gcd(m_2, m_3) + \gcd(m_4, m_5)m_1.$$

Set $J = (g_1, g_2, g_3)$. Note that $\sqrt{x_{16} \gcd(m_1, m_3) \gcd(m_2, m_3)} = m_3$, where \sqrt{m} denotes a product of all variables those are appear in a monomial m . Then it is easy to see that $\sqrt{J} \subset I$. We prove the opposite inclusion.

Since $x_{16}m_4m_5 = x_{16}(g_2 - f_1m_1)(g_3 - f_2m_2) \in J$, we have $x_7x_{14}x_{15}x_{17}x_{20}m_4 \in \sqrt{J}$ and $x_4x_{11}x_{12}x_{18}x_{19}m_5 \in \sqrt{J}$. Then $x_{17}m_4^2 = x_{17}m_4(g_2 - f_1m_1) \in \sqrt{J}$ because $x_7x_{14}x_{15}x_{20}$ divides m_1 . Therefore we have $x_{17}m_4 \in \sqrt{J}$, and thus $x_{17}f_1m_1 = x_{17}(g_2 - m_4) \in \sqrt{J}$. Since x_{17} divides $\gcd(m_2, m_3)$, the belonging $\gcd(m_2, m_3)g_1 \in J$ implies $x_{16} \gcd(m_2, m_3)f_1 \in \sqrt{J}$. Then $x_{16} \gcd(m_2, m_3) \gcd(m_1, m_3) \in \sqrt{J}$ and $x_{16} \gcd(m_2, m_3) \gcd(m_4, m_5)m_2 \in \sqrt{J}$ because m_4 divides $\gcd(m_1, m_3) \gcd(m_4, m_5)m_2$. Hence $m_3, \gcd(m_4, m_5)m_2 \in \sqrt{J}$. Similarly, we have $\gcd(m_4, m_5)m_1 \in \sqrt{J}$. Then $g_2 \in J$ (resp. $g_3 \in J$) implies that $m_1, m_4 \in \sqrt{J}$ (resp. $m_2, m_5 \in \sqrt{J}$), as desired.

Case (2): We prove that the following 3 elements g_1, g_2, g_3 generate I up to radical:

$$g_1 = x_{13} \left(\frac{m_4}{x_{13}} + m_3 \right), \quad g_2 = m_1m_2 \left(\frac{m_4}{x_{13}} + m_3 \right) + m_5, \quad g_3 = m_1 + m_2 + m_3.$$

Set $J = (g_1, g_2, g_3)$. Then it is clear that $\sqrt{J} \subset I$. We prove the opposite inclusion.

Since $x_{13}m_5 = x_{13}(g_2 - m_1m_2((m_4/x_{13}) + m_3)) = x_{13}g_2 - m_1m_2g_1 \in J$, we have $m_5 \in \sqrt{J}$. Then $x_{13}x_{18}m_3^2 = x_{18}m_3(g_1 - m_4) \in \sqrt{J}$ because $x_{18}m_3m_4$ is divisible by $m_5 \in \sqrt{J}$. This implies $x_{18}m_3 \in \sqrt{J}$. Since x_{18} is a common factor of m_1 and m_2 , we have $m_3^2 = m_3(g_3 - m_1 - m_2) \in \sqrt{J}$. That is $m_3 \in \sqrt{J}$, and thus $m_4 = g_1 - x_{13}m_3, m_1 + m_2 = g_3 - m_3, m_1m_2(m_4/x_{13}) = g_2 - m_1m_2m_3 - m_5 \in \sqrt{J}$.

Since $m_1 m_2$ is divisible by m_4/x_{13} , we have $m_1 m_2 \in \sqrt{J}$. Therefore $m_1, m_2 \in \sqrt{J}$, as desired.

Case (3): Since the length of the Lyubeznik resolution ([18]) of $I = (m_5, m_1, m_2, m_3, m_4)$ with respect to this order of generators is equal to 3, the inequality $\text{ara } I \leq 3$ follows from [11]. \square

8. THE CASE $\text{arithdeg } I \leq 4$

Let I be a squarefree monomial ideal of S with $\text{arithdeg } I \leq 4$. In this section, we prove that $\text{ara } I = \text{pd}_S S/I$ holds for such an ideal I .

Theorem 8.1. *Let S be a polynomial ring over a field K and I a squarefree monomial ideal of S with $\text{arithdeg } I \leq 4$. Then*

$$\text{ara } I = \text{pd}_S S/I.$$

To prove the theorem, first, we reduce to simple cases. Then we find $\text{pd}_S S/I$ elements which generate I up to radical. On this step, we use the result due to Schmitt and Vogel [20].

Lemma 8.2 (Schmitt and Vogel [20, Lemma, p. 249]). *Let R be a commutative ring and \mathcal{P} a finite subset of R . Suppose that subsets $\mathcal{P}_0, \mathcal{P}_1, \dots, \mathcal{P}_r$ of \mathcal{P} satisfy the following three conditions:*

$$(SV1) \bigcup_{\ell=0}^r \mathcal{P}_\ell = \mathcal{P};$$

$$(SV2) |\mathcal{P}_0| = 1;$$

(SV3) *for all $\ell > 0$ and for all elements $a, a'' \in \mathcal{P}_\ell$ with $a \neq a''$, there exist an integer $\ell' < \ell$ and an element $a' \in \mathcal{P}_{\ell'}$ such that $aa'' \in (a')$.*

Set $I = (\mathcal{P})$ and

$$g_\ell = \sum_{a \in \mathcal{P}_\ell} a, \quad \ell = 0, 1, \dots, r.$$

Then g_0, g_1, \dots, g_r generate I up to radical.

First, we reduce to the case where $\dim \mathcal{H}(I^*) \leq 1$. To do this, we need the following lemma.

Lemma 8.3. *Let $I = I_\Delta$ be a squarefree monomial ideal of $S = K[x_1, \dots, x_n]$ with $\text{arithdeg } I \leq 4$ and $\text{indeg } I \geq 2$. Then there exists a squarefree monomial ideal $I' = I_{\Delta'}$ such that $\dim \mathcal{H}(I'^*) \leq 1$ and that Δ is obtained from Δ' adding cones recursively.*

Proof. Since $\mu(I^*) \leq 4$ and $\text{height } I^* \geq 2$, we have $\dim \mathcal{H}(I^*) \leq 2$ by [14, Proposition 3.4]. We prove the lemma by induction on the number c of 2-faces of $\mathcal{H}(I^*)$.

If $c = 0$, then $\dim \mathcal{H}(I^*) \leq 1$ and we may take $\Delta' = \Delta$.

Assume $c \geq 1$. Then $\text{arithdeg } I = 4$ since $\text{indeg } I \geq 2$, and there exists a 2-face $F \in \mathcal{H}(I^*)$. We may assume $F = \{1, 2, 3\}$ and let x_1, \dots, x_t ($t \geq 1$) be all defining variables of F . Let

$$I = I_\Delta = P_1 \cap P_2 \cap P_3 \cap P_4$$

be the minimal prime decomposition of I . Then we can write $P_i = P'_i + (x_1)$ for $i = 1, 2, 3$, where $x_1 \notin P'_i$. Note that $x_1 \notin P_4$. Set

$$(8.1) \quad I' = P'_1 \cap P'_2 \cap P'_3 \cap P_4 \quad (\subset S' = K[x_2, \dots, x_n]),$$

and let Δ' be the simplicial complex associated with I' . Then $\Delta = \Delta' \cup \text{cone}_{x_1}(G_4)$, where G_4 is the facet of Δ' corresponding to P_4 . If $t > 1$, then $\mathcal{H}(I'^*) = \mathcal{H}(I^*)$

and (8.1) is still the minimal prime decomposition of I' . Hence we can reduce the case $t = 1$ by applying the above operation $t - 1$ times. If (8.1) is not a minimal prime decomposition of I' , then $I' = P'_1 \cap P'_2 \cap P'_3$ and $\dim \mathcal{H}(I'^*) \leq 1$. Otherwise $\mathcal{H}(I'^*) = \mathcal{H}(I^*) \setminus \{F\}$ and the number of 2-feces of $\mathcal{H}(I'^*)$ is smaller than that of $\mathcal{H}(I^*)$, as required. \square

Barile and Terai [5, Theorem 2, p. 3694] (see also [12, Section 5]) proved that if $\text{ara } I' = \text{pd}_S S'/I'$ holds, then $\text{ara } I = \text{pd}_S S/I$ also holds, where the notations are the same as in Lemma 8.3. Therefore we only need to consider the case where $\dim \mathcal{H}(I^*) \leq 1$.

Next, we reduce to the case where $\mathcal{H}(I^*)$ is connected.

Lemma 8.4. *Let I_1, I_2 be squarefree monomial ideals of S . Suppose that $X(I_1) \cap X(I_2) = \emptyset$. Then*

$$\text{ara}(I_1 \cap I_2) \leq \text{ara } I_1 + \text{ara } I_2 - 1.$$

Moreover, if $\text{ara } I_i = \text{pd}_S S/I_i$ holds for $i = 1, 2$, then $\text{ara}(I_1 \cap I_2) = \text{pd}_S S/(I_1 \cap I_2)$ also holds.

Proof. Let $\text{ara } I_i = s_i + 1$ for $i = 1, 2$. Then there exist $s_1 + 1$ elements $f_0, f_1, \dots, f_{s_1} \in S$ (resp. $s_2 + 1$ elements $g_0, g_1, \dots, g_{s_2} \in S$) such that those generate I_1 (resp. I_2) up to radical.

Set

$$h_\ell = \sum_{j=0}^{\ell} f_{\ell-j} g_j, \quad \ell = 0, 1, \dots, s_1 + s_2$$

and $J = (h_0, h_1, \dots, h_{s_1+s_2})$, where $f_i = 0$ (resp. $g_j = 0$) if $i > s_1$ (resp. $j > s_2$). If $\sqrt{J} = I_1 \cap I_2$, then

$$\text{ara}(I_1 \cap I_2) \leq s_1 + s_2 + 1 = \text{ara } I_1 + \text{ara } I_2 - 1.$$

We prove $\sqrt{J} = I_1 \cap I_2$. Since $h_\ell \in I_1 \cap I_2$, we have $\sqrt{J} \subset I_1 \cap I_2$. We prove the opposite inclusion. Note that $I_1 \cap I_2 = I_1 I_2$ since $X(I_1) \cap X(I_2) = \emptyset$. By Lemma 8.2, we have $f_i g_j \in \sqrt{J}$ for all $0 \leq i \leq s_1$ and for all $0 \leq j \leq s_2$. Then

$$(f_0, f_1, \dots, f_{s_1})(g_0, g_1, \dots, g_{s_2}) \subset \sqrt{J}.$$

Hence

$$\sqrt{(f_0, f_1, \dots, f_{s_1})(g_0, g_1, \dots, g_{s_2})} \subset \sqrt{J}.$$

On the other hand,

$$\begin{aligned} \sqrt{(f_0, f_1, \dots, f_{s_1})(g_0, g_1, \dots, g_{s_2})} &\supset \sqrt{(f_0, f_1, \dots, f_{s_1})} \sqrt{(g_0, g_1, \dots, g_{s_2})} \\ &= I_1 I_2 = I_1 \cap I_2. \end{aligned}$$

Therefore we have $I_1 \cap I_2 \subset \sqrt{J}$, as desired.

To prove the second part of the lemma, we set $I = I_1 \cap I_2$. Then $I^* = I_1^* + I_2^*$. Note that $X(I_1) \cap X(I_2) = \emptyset$ implies $X(I_1^*) \cap X(I_2^*) = \emptyset$. Since we have already known the inequality $\text{ara } I \geq \text{pd}_S S/I$ by (1.1), it is sufficient to prove the opposite inequality.

By assumption, we have

$$(8.2) \quad \text{ara } I = \text{ara}(I_1 \cap I_2) \leq \text{ara } I_1 + \text{ara } I_2 - 1 = \text{pd}_S S/I_1 + \text{pd}_S S/I_2 - 1.$$

On the other hand, since $X(I_1^*) \cap X(I_2^*) = \emptyset$, we have

$$\beta_{i,j}(S/I^*) = \sum_{\ell=0}^j \sum_{m=0}^i \beta_{m,\ell}(S/I_1^*) \beta_{i-m,j-\ell}(S/I_2^*).$$

Assume that the regularity of S/I_1^* (resp. S/I_2^*) is given by $\beta_{i_1,j_1}(S/I_1^*) \neq 0$ (resp. $\beta_{i_2,j_2}(S/I_2^*) \neq 0$). Then

$$\text{reg } S/I_1^* = j_1 - i_1, \quad \text{reg } S/I_2^* = j_2 - i_2,$$

and $\beta_{i_1+i_2,j_1+j_2}(S/I^*) \geq \beta_{i_1,j_1}(S/I_1^*) \beta_{i_2,j_2}(S/I_2^*) > 0$. Hence,


$$(8.3) \quad \text{reg } I^* = \text{reg}(S/I^*) + 1 \geq j_1 + j_2 - (i_1 + i_2) + 1.$$


Therefore

$$\begin{aligned} \text{pd}_S S/I &= \text{reg } I^* \geq \text{reg } S/I_1^* + \text{reg } S/I_2^* + 1 && \text{(by (8.3))} \\ &= \text{reg } I_1^* + \text{reg } I_2^* - 1 \\ &= \text{pd}_S S/I_1 + \text{pd}_S S/I_2 - 1 \\ &\geq \text{ara } I && \text{(by (8.2)).} \end{aligned}$$


□

Let I be a squarefree monomial ideal with $\text{arithdeg } I \leq 4$. By Remark 5.5 we may assume $\text{indeg } I \geq 2$. It was proved in [14, Theorems 5.1 and 6.1] that $\text{ara } I = \text{pd}_S S/I$ holds when $\text{arithdeg } I - \text{indeg } I \leq 1$. Hence we only need to consider the case where $\text{arithdeg } I = 4$ and $\text{indeg } I = 2$, equivalently, $\mu(I^*) = 4$ and $\text{height } I^* = 2$. Then corresponding hypergraphs $\mathcal{H} := \mathcal{H}(I^*)$ were classified in [15, Section 3]. As a consequence of Lemmas 8.3 and 8.4, we may assume that \mathcal{H} is connected and $\dim \mathcal{H} \leq 1$. Moreover since we know $\text{ara } I = \text{pd}_S S/I$ holds when $\text{arithdeg } I = \text{reg } I$, i.e., $\mu(I^*) = \text{pd}_S S/I^*$ by [14, Theorem 5.1], we may assume that $W(\mathcal{H}) \neq \emptyset$ by Proposition 4.1. Thus \mathcal{H} coincides with one of the following hypergraphs:





Note that \mathcal{H} is contained in the hypergraph \mathcal{H}_{17} :



Throughout, we label the vertices of \mathcal{H} as above. Then I is of the form $I = P_1 \cap P_2 \cap P_3 \cap P_4$ with

$$\begin{aligned} P_1 &= (x_{11}, \dots, x_{1i_1}, x_{41}, \dots, x_{4i_4}, x_{51}, \dots, x_{5i_5}), \\ P_2 &= (x_{11}, \dots, x_{1i_1}, x_{21}, \dots, x_{2i_2}, x_{61}, \dots, x_{6i_6}, y_{21}, \dots, y_{2j_2}), \\ P_3 &= (x_{21}, \dots, x_{2i_2}, x_{31}, \dots, x_{3i_3}, x_{51}, \dots, x_{5i_5}, y_{31}, \dots, y_{3j_3}), \\ P_4 &= (x_{31}, \dots, x_{3i_3}, x_{41}, \dots, x_{4i_4}, x_{61}, \dots, x_{6i_6}, y_{41}, \dots, y_{4j_4}), \end{aligned}$$

where $\{x_{st}\}, \{y_{uv}\}$ are all distinct variables of S and $i_s \geq 0, j_u \geq 0$. Then we have

$$I^* = (X_1 X_4 X_5, X_1 X_2 X_6 Y_2, X_2 X_3 X_5 Y_3, X_3 X_4 X_6 Y_4),$$

where

$$X_s = x_{s1} \cdots x_{si_s}, \quad s = 1, 2, 3, 4; \quad Y_u = y_{u1} \cdots y_{uj_u}, \quad u = 2, 3, 4.$$

Here we set $X_s = 1$ (resp. $Y_u = 1$) when $i_s = 0$ (resp. $j_u = 0$).

Let $N = i_1 + \cdots + i_6 + j_2 + j_3 + j_4$. Then one can easily construct a graded minimal free resolution of I^* and compute $\text{reg } I^*$.

Lemma 8.5. *Let I be a squarefree monomial ideal with $\text{indeg } I^* \geq 2$. Suppose that $\mathcal{H} := \mathcal{H}(I^*)$ coincides with one of $\mathcal{H}_1, \dots, \mathcal{H}_{24}$.*

- (1) *When \mathcal{H} coincides with one of $\mathcal{H}_{11}, \mathcal{H}_{17}, \mathcal{H}_{20}$,*

$$\text{pd}_S S/I = \text{reg } I^* = \max\{N - j_2 - 2, N - j_3 - 2, N - j_4 - 2\}.$$

- (2) *When $\mathcal{H} = \mathcal{H}_{22}$,*

$$\text{pd}_S S/I = \text{reg } I^* = \max\{N - j_2 - 2, N - j_3 - 2\}.$$

- (3) *When \mathcal{H} coincides with one of $\mathcal{H}_4, \mathcal{H}_5, \mathcal{H}_{12}, \mathcal{H}_{13}, \mathcal{H}_{24}$,*

$$\text{pd}_S S/I = \text{reg } I^* = \max\{N - j_2 - 2, N - j_4 - 2\}.$$

- (4) *Otherwise, $\text{pd}_S S/I = \text{reg } I^* = N - 2$.*

Now we find $\text{pd}_S S/I$ elements of I which generate I up to radical. First, we consider the case of \mathcal{H}_{17} . The following construction for $\mathcal{H} = \mathcal{H}_{17}$ is also valid for the other cases except for the two cases of \mathcal{H}_1 and \mathcal{H}_{14} .

Set

$$r_1 = N - j_4 - 3, \quad r_2 = N - j_3 - 3, \quad r_3 = N - j_2 - 3, \quad r = \max\{r_1, r_2, r_3\}.$$

Then $\text{pd}_S S/I = r + 1$. We define sets $\mathcal{P}_\ell^{(1)}, \dots, \mathcal{P}_\ell^{(8)}$ as follows and set $\mathcal{P}_\ell = \mathcal{P}_\ell^{(1)} \cup \dots \cup \mathcal{P}_\ell^{(8)}$ for $\ell = 0, 1, \dots, r$:

$$\begin{aligned} \mathcal{P}_\ell^{(1)} &= \{x_{1\ell_1}x_{3\ell_3} : \ell_1 + \ell_3 = \ell + 2; 1 \leq \ell_s \leq i_s (s = 1, 3)\}, \\ \mathcal{P}_\ell^{(2)} &= \left\{ x_{1\ell_1}w_{3\ell_3}w_{4\ell_4} : \begin{array}{l} \ell_1 + \ell_3 + \ell_4 + i_3 = \ell + 3, \\ 1 \leq \ell_1 \leq i_1; 1 \leq \ell_3 \leq i_2 + i_5 + j_3; 1 \leq \ell_4 \leq i_4 + i_6 + j_4 \end{array} \right\}, \\ \mathcal{P}_\ell^{(3)} &= \left\{ x_{3\ell_3}w_{1\ell_1}w_{2\ell_2} : \begin{array}{l} \ell_3 + \ell_1 + \ell_2 + i_1 = \ell + 3, \\ 1 \leq \ell_3 \leq i_3; 1 \leq \ell_1 \leq i_4 + i_5; 1 \leq \ell_2 \leq i_2 + i_6 + j_2 \end{array} \right\}, \\ \mathcal{P}_\ell^{(4)} &= \{x_{2\ell_2}x_{4\ell_4} : \ell_2 + \ell_4 + i_1 + i_3 = \ell + 2; 1 \leq \ell_s \leq i_s (s = 2, 4)\}, \\ \mathcal{P}_\ell^{(5)} &= \left\{ x_{4\ell_4}w_{2\ell_2}w_{3\ell_3} : \begin{array}{l} \ell_4 + (\ell_2 - i_2) + (\ell_3 - i_2) + i_1 + i_2 + i_3 = \ell + 3, \\ 1 \leq \ell_4 \leq i_4; i_2 < \ell_2 \leq i_2 + i_6 + j_2; i_2 < \ell_3 \leq i_2 + i_5 + j_3 \end{array} \right\}, \\ \mathcal{P}_\ell^{(6)} &= \left\{ x_{2\ell_2}x_{5\ell_5}w_{4\ell_4} : \begin{array}{l} \ell_2 + \ell_5 + (\ell_4 - i_4) + i_1 + i_3 + i_4 = \ell + 3, \\ 1 \leq \ell_s \leq i_s (s = 2, 5); i_4 < \ell_4 \leq i_4 + i_6 + j_4 \end{array} \right\}, \\ \mathcal{P}_\ell^{(7)} &= \{x_{5\ell_5}x_{6\ell_6} : \ell_5 + \ell_6 + i_1 + \dots + i_4 = \ell + 2; 1 \leq \ell_s \leq i_s (s = 5, 6)\}, \\ \mathcal{P}_\ell^{(8)} &= \left\{ x_{5\ell_5}y_{2\ell_2}y_{4\ell_4} : \begin{array}{l} \ell_5 + \ell_2 + \ell_4 + i_1 + \dots + i_4 + i_6 = \ell + 3, \\ 1 \leq \ell_5 \leq i_5; 1 \leq \ell_u \leq j_u (u = 2, 4) \end{array} \right\}. \end{aligned}$$

Here,

$$\begin{aligned} w_{1\ell_1} &= \begin{cases} x_{4\ell_1}, & 1 \leq \ell_1 \leq i_4, \\ x_{5\ell_1 - i_4}, & i_4 < \ell_1 \leq i_4 + i_5, \end{cases} \\ w_{2\ell_2} &= \begin{cases} x_{2\ell_2}, & 1 \leq \ell_2 \leq i_2, \\ x_{6\ell_2 - i_2}, & i_2 < \ell_2 \leq i_2 + i_6, \\ y_{2\ell_2 - i_2 - i_6}, & i_2 + i_6 < \ell_2 \leq i_2 + i_6 + j_2, \end{cases} \\ w_{3\ell_3} &= \begin{cases} x_{2\ell_3}, & 1 \leq \ell_3 \leq i_2, \\ x_{5\ell_3 - i_2}, & i_2 < \ell_3 \leq i_2 + i_5, \\ y_{3\ell_3 - i_2 - i_5}, & i_2 + i_5 < \ell_3 \leq i_2 + i_5 + j_3, \end{cases} \\ w_{4\ell_4} &= \begin{cases} x_{4\ell_4}, & 1 \leq \ell_4 \leq i_4, \\ x_{6\ell_4 - i_4}, & i_4 < \ell_4 \leq i_4 + i_6, \\ y_{4\ell_4 - i_4 - i_6}, & i_4 + i_6 < \ell_4 \leq i_4 + i_6 + j_4. \end{cases} \end{aligned}$$

Note that for each k ($k = 1, 2, \dots, 8$), the range of ℓ with $\mathcal{P}_\ell^{(k)} \neq \emptyset$ is given by the following list:


$$\begin{aligned} \mathcal{P}_\ell^{(1)} &: 0 \leq \ell \leq i_1 + i_3 - 2, \\ \mathcal{P}_\ell^{(2)} &: i_3 \leq \ell \leq N - j_2 - 3 = r_3, \\ \mathcal{P}_\ell^{(3)} &: i_1 \leq \ell \leq N - j_3 - j_4 - 3, \\ \mathcal{P}_\ell^{(4)} &: i_1 + i_3 \leq \ell \leq i_1 + \dots + i_4 - 2, \\ \mathcal{P}_\ell^{(5)} &: i_1 + i_2 + i_3 \leq \ell \leq N - j_4 - 3 = r_1, \\ \mathcal{P}_\ell^{(6)} &: i_1 + i_3 + i_4 \leq \ell \leq N - j_2 - j_3 - 3, \\ \mathcal{P}_\ell^{(7)} &: i_1 + \dots + i_4 \leq \ell \leq i_1 + \dots + i_6 - 2, \\ \mathcal{P}_\ell^{(8)} &: i_1 + \dots + i_4 + i_6 \leq \ell \leq N - j_3 - 3 = r_2. \end{aligned}$$

Now, we verify that \mathcal{P}_ℓ , $\ell = 0, 1, \dots, r$ satisfy the conditions (SV1), (SV2), and (SV3). In this case, the condition (SV1) means that $\bigcup_{\ell=0}^r \mathcal{P}_\ell$ generates I , and it is satisfied. Since $\mathcal{P}_0 = \mathcal{P}_0^{(1)} = \{x_{11}x_{31}\}$, the condition (SV2) is also satisfied. To check the condition (SV3), let a, a'' be two distinct elements in \mathcal{P}_ℓ . We denote the indices of a (resp. a'') by ℓ_s (resp. ℓ''_s). First suppose $a, a'' \in \mathcal{P}_\ell^{(k)}$. Then there exists s such that $\ell_s \neq \ell''_s$ and we may assume $\ell_s < \ell''_s$. Replacing ℓ''_s to ℓ_s in a'' , we obtain required elements $a' \in \mathcal{P}_{\ell'}^{(k)}$ with $\ell' < \ell$. For example, take two distinct elements $a = x_{1\ell_1}x_{3\ell_3}$, $a'' = x_{1\ell''_1}x_{3\ell''_3} \in \mathcal{P}_\ell^{(1)}$ with $\ell_1 < \ell''_1$. Then $a' = x_{1\ell_1}x_{3\ell''_3} \in \mathcal{P}_{\ell'}^{(1)}$, where $\ell' = \ell_1 + \ell''_3 - 2 < \ell''_1 + \ell''_3 - 2 = \ell$. Next, we assume $a \in \mathcal{P}_\ell^{(k)}$ and $a'' \in \mathcal{P}_\ell^{(k')}$ with $k < k'$. If $k = 1$, $k' = 2$, then $a = x_{1\ell_1}x_{3\ell_3}$, $a'' = x_{1\ell''_1}w_{3\ell''_3}w_{4\ell''_4}$. We can take $a' = x_{1\ell''_1}x_{3\ell_3} \in \mathcal{P}_{\ell'}^{(1)}$, where

$$\begin{aligned} \ell' &= \ell''_1 + \ell_3 - 2 \\ &= (\ell + 3 - (\ell''_3 + \ell''_4 + i_3)) + \ell_3 - 2 \\ &= \ell + 1 - \ell''_3 - \ell''_4 - (i_3 - \ell_3) \\ &< \ell. \end{aligned}$$

Similarly, we can check the existence of required ℓ' and $a' \in \mathcal{P}_{\ell'}^{(k')}$ for other pairs (k, k') ; we just mention the choices for a' in Table 1.

Next we consider the two exceptional cases: $\mathcal{H} = \mathcal{H}_1, \mathcal{H}_{14}$:



The case of \mathcal{H}_1 is easy because in this case,

$$\begin{aligned} I &= (x_{11}, \dots, x_{1i_1}, x_{41}, \dots, x_{4i_4}) \cap (x_{11}, \dots, x_{1i_1}, x_{21}, \dots, x_{2i_2}) \\ &\quad \cap (x_{21}, \dots, x_{2i_2}, x_{31}, \dots, x_{3i_3}) \cap (x_{31}, \dots, x_{3i_3}, x_{41}, \dots, x_{4i_4}) \\ &= (x_{1\ell_1}x_{3\ell_3} : 1 \leq \ell_s \leq i_s \ (s = 1, 3)) + (x_{2\ell_2}x_{4\ell_4} : 1 \leq \ell_s \leq i_s \ (s = 2, 4)) \\ &= (x_{11}, \dots, x_{1i_1}) \cap (x_{31}, \dots, x_{3i_3}) + (x_{21}, \dots, x_{2i_2}) \cap (x_{41}, \dots, x_{4i_4}). \end{aligned}$$

TABLE 1. The choices for a' in the case of \mathcal{H}_{17} .

k	k''	k'	a'
1	2	1	$x_{1\ell_1'} x_{3\ell_3}$
1	3	1	$x_{1\ell_1} x_{3\ell_3''}$
1	4-8		These cases do not occur.
2	3	1	$x_{1\ell_1} x_{3\ell_3''}$
2	4	2	$x_{1\ell_1} x_{2\ell_2''} x_{4\ell_4''} = x_{1\ell_1} w_{3\ell_2''} w_{4\ell_4''}$
2	5	2	$x_{1\ell_1} w_{3\ell_3''} x_{4\ell_4''} = x_{1\ell_1} w_{3\ell_3''} w_{4\ell_4''}$
2	6	2	$x_{1\ell_1} x_{2\ell_2''} w_{4\ell_4''} = x_{1\ell_1} w_{3\ell_2''} w_{4\ell_4''}$
2	7	2	$x_{1\ell_1} x_{5\ell_5''} x_{6\ell_6''} = x_{1\ell_1} w_{3i_2+\ell_5''} w_{4i_4+\ell_6''}$
2	8	2	$x_{1\ell_1} x_{5\ell_5''} y_{4\ell_4''} = x_{1\ell_1} w_{3i_2+\ell_5''} w_{4i_4+i_6+\ell_4''}$
3	4	3	$x_{3\ell_3} x_{4\ell_4''} x_{2\ell_2''} = x_{3\ell_3} w_{1\ell_4''} w_{2\ell_2''}$
3	5	3	$x_{3\ell_3} x_{4\ell_4''} w_{2\ell_2''} = x_{3\ell_3} w_{1\ell_4''} w_{2\ell_2''}$
3	6	3	$x_{3\ell_3} x_{5\ell_5''} x_{2\ell_2''} = x_{3\ell_3} w_{1i_4+\ell_5''} w_{2\ell_2''}$
3	7	3	$x_{3\ell_3} x_{5\ell_5''} x_{6\ell_6''} = x_{3\ell_3} w_{1i_4+\ell_5''} w_{2i_2+\ell_6''}$
3	8	3	$x_{3\ell_3} x_{5\ell_5''} y_{2\ell_2''} = x_{3\ell_3} w_{1i_4+\ell_5''} w_{2i_2+i_6+\ell_2''}$
4	5	4	$x_{2\ell_2} x_{4\ell_4''}$
4	6	4	$x_{2\ell_2''} x_{4\ell_4}$
4	7, 8		These cases do not occur.
5	6	4	$x_{2\ell_2''} x_{4\ell_4}$
5	7	5	$x_{4\ell_4} x_{6\ell_6''} x_{5\ell_5''} = x_{4\ell_4} w_{2i_2+\ell_6''} w_{3i_2+\ell_5''}$
5	8	5	$x_{4\ell_4} y_{2\ell_2''} x_{5\ell_5''} = x_{4\ell_4} w_{2i_2+i_6+\ell_2''} w_{3i_2+\ell_5''}$
6	7	6	$x_{2\ell_2} x_{5\ell_5''} x_{6\ell_6''} = x_{2\ell_2} x_{5\ell_5''} w_{4i_4+\ell_6''}$
6	8	6	$x_{2\ell_2} x_{5\ell_5''} y_{4\ell_4''} = x_{2\ell_2} x_{5\ell_5''} w_{4i_4+i_6+\ell_4''}$
7	8	7	$x_{5\ell_5''} x_{6\ell_6}$

Hence, I is the sum of the two squarefree monomial ideals $I_1 := (x_{11}, \dots, x_{1i_1}) \cap (x_{31}, \dots, x_{3i_3})$ and $I_2 := (x_{21}, \dots, x_{2i_2}) \cap (x_{41}, \dots, x_{4i_4})$ with $\text{arithdeg } I_i = \text{indeg } I_i$ ($i = 1, 2$). It is known by Schmitt and Vogel [20, Theorem 1, p. 247] that $\text{ara } I_i = \text{pd}_S S/I_i$ ($i = 1, 2$). Since $X(I_1) \cap X(I_2) = \emptyset$, we have $\text{ara } I = \text{pd}_S S/I$ by Proposition 5.4.

For the case of \mathcal{H}_{14} , we use Lemma 8.2 again. In this case, $\text{pd}_S S/I = N - 2$. Note that $N = i_1 + \dots + i_6$ since $j_2 = j_3 = j_4 = 0$. We define $\mathcal{P}_\ell := \mathcal{P}_\ell^{(1)} \cup \dots \cup \mathcal{P}_\ell^{(5)}$, $\ell = 0, 1, \dots, N - 3$, by

$$\begin{aligned}
\mathcal{P}_\ell^{(1)} &= \{x_{1\ell_1} x_{3\ell_3} : \ell_1 + \ell_3 = \ell + 2; 1 \leq \ell_s \leq i_s (s = 1, 3)\}, \\
\mathcal{P}_\ell^{(2)} &= \left\{ x_{1\ell_1} w_{3\ell_3} w_{4\ell_4} : \begin{array}{l} \ell_1 + \ell_3 + \ell_4 + i_3 = \ell + 3, \\ 1 \leq \ell_1 \leq i_1; 1 \leq \ell_3 \leq i_2 + i_5; 1 \leq \ell_4 \leq i_4 + i_6, \\ \ell_3 \leq i_2 \text{ or } \ell_4 \leq i_4 \end{array} \right\}, \\
\mathcal{P}_\ell^{(3)} &= \left\{ x_{3\ell_3} w_{1\ell_1} w_{2\ell_2} : \begin{array}{l} \ell_3 + \ell_1 + \ell_2 + i_1 = \ell + 3, \\ 1 \leq \ell_3 \leq i_3; 1 \leq \ell_1 \leq i_4 + i_5; 1 \leq \ell_2 \leq i_2 + i_6, \\ \ell_1 \leq i_4 \text{ or } \ell_2 \leq i_2 \end{array} \right\}, \\
\mathcal{P}_\ell^{(4)} &= \{x_{2\ell_2} x_{4\ell_4} : \ell_2 + \ell_4 + i_1 + i_3 = \ell + 2; 1 \leq \ell_s \leq i_s (s = 2, 4)\}, \\
\mathcal{P}_\ell^{(5)} &= \{x_{5\ell_5} x_{6\ell_6} : \ell_5 + \ell_6 + i_1 + \dots + i_4 = \ell + 3; 1 \leq \ell_s \leq i_s (s = 5, 6)\}.
\end{aligned}$$

TABLE 2. The choices for a' in the case of \mathcal{H}_{14} .

k	k''	k'	additional condition	a'
1	2	1		$x_{1\ell_1} x_{3\ell_3}$
1	3	1		$x_{1\ell_1} x_{3\ell_3}$
1	4, 5		These cases do not occur.	
2	3	1		$x_{1\ell_1} x_{3\ell_3}$
2	4	2		$x_{1\ell_1} x_{2\ell_2} x_{4\ell_4} = x_{1\ell_1} w_{3\ell_2} w_{4\ell_4}$
2	5	2	$\ell_3 \leq i_2$	$x_{1\ell_1} w_{3\ell_3} x_{6\ell_6} = x_{1\ell_1} w_{3\ell_3} w_{4i_4 + \ell_6}$
2	5	2	$\ell_4 \leq i_4$	$x_{1\ell_1} x_{5\ell_5} w_{4\ell_4} = x_{1\ell_1} w_{3i_2 + \ell_5} w_{4\ell_4}$
3	4	3		$x_{3\ell_3} x_{4\ell_4} x_{2\ell_2} = x_{3\ell_3} w_{1\ell_4} w_{2\ell_2}$
3	5	3	$\ell_1 \leq i_4$	$x_{3\ell_3} w_{1\ell_1} x_{6\ell_6} = x_{3\ell_3} w_{1\ell_1} w_{2i_2 + \ell_6}$
3	5	3	$\ell_2 \leq i_2$	$x_{3\ell_3} x_{5\ell_5} w_{2\ell_2} = x_{3\ell_3} w_{1i_4 + \ell_5} w_{2\ell_2}$
4	5		This case does not occur.	

Here,

$$w_{1\ell_1} = \begin{cases} x_{4\ell_1}, & 1 \leq \ell_1 \leq i_4, \\ x_{5\ell_1 - i_4}, & i_4 < \ell_1 \leq i_4 + i_5, \end{cases} \quad w_{2\ell_2} = \begin{cases} x_{2\ell_2}, & 1 \leq \ell_2 \leq i_2, \\ x_{6\ell_2 - i_2}, & i_2 < \ell_2 \leq i_2 + i_6, \end{cases}$$

$$w_{3\ell_3} = \begin{cases} x_{2\ell_3}, & 1 \leq \ell_3 \leq i_2, \\ x_{5\ell_3 - i_2}, & i_2 < \ell_3 \leq i_2 + i_5, \end{cases} \quad w_{4\ell_4} = \begin{cases} x_{4\ell_4}, & 1 \leq \ell_4 \leq i_4, \\ x_{6\ell_4 - i_4}, & i_4 < \ell_4 \leq i_4 + i_6. \end{cases}$$

Note that for each k ($k = 1, 2, \dots, 5$), the range of ℓ with $\mathcal{P}_\ell^{(k)} \neq \emptyset$ is given by the following list:

$$\begin{aligned} \mathcal{P}_\ell^{(1)} &: 0 \leq \ell \leq i_1 + i_3 - 2, \\ \mathcal{P}_\ell^{(2)} &: i_3 \leq \ell \leq \max\{N - i_5 - 3, N - i_6 - 3\}, \\ \mathcal{P}_\ell^{(3)} &: i_1 \leq \ell \leq \max\{N - i_5 - 3, N - i_6 - 3\}, \\ \mathcal{P}_\ell^{(4)} &: i_1 + i_3 \leq \ell \leq i_1 + \dots + i_4 - 2, \\ \mathcal{P}_\ell^{(5)} &: i_1 + \dots + i_4 - 1 \leq \ell \leq i_1 + \dots + i_6 - 3 = N - 3. \end{aligned}$$

By a similar way to the case of \mathcal{H}_{17} , we can check that \mathcal{P}_ℓ , $\ell = 0, 1, \dots, N - 3$ satisfy the conditions (SV1), (SV2), and (SV3). For the condition (SV3), as in \mathcal{H}_{17} , we list the choices for $a' \in \mathcal{P}_\ell^{(k')}$ from given $a, a'' \in \mathcal{P}_\ell$ in Table 2.

This completes the proof of Theorem 8.1.

Remark 8.6. For the exceptional case $\mathcal{H} = \mathcal{H}_{14}$, we can also consider the corresponding elements on the construction for \mathcal{H}_{17} ; in fact, these generate I up to radical. But in this case, the range of ℓ with $\mathcal{P}_\ell \neq \emptyset$ is $0 \leq \ell \leq N - 2$ because of $\mathcal{P}_\ell^{(7)}$. Therefore, we cannot obtain $\text{ara } I$ from this construction. This is also true for $\mathcal{H} = \mathcal{H}_1$.

Acknowledgment. The authors thank the referee for reading the manuscript carefully.

REFERENCES

- [1] M. Barile, *On the number of equations defining certain varieties*, *manuscripta math.* **91** (1996), 483–494.

- [2] M. Barile, *Arithmetical ranks of Stanley–Reisner ideals via linear algebra*, *Comm. Algebra* **36** (2008), 4540–4556.
- [3] M. Barile, *On the arithmetical rank of the edge ideals of forests*, *Comm. Algebra* **36** (2008), 4678–4703.
- [4] M. Barile, D. Kiani, F. Mohammadi and S. Yassemi, *Arithmetical rank of the cyclic and bicyclic graphs*, to appear in *J. Pure Appl. Algebra*, arXiv:0805.1657.
- [5] M. Barile and N. Terai, *Arithmetical ranks of Stanley–Reisner ideals of simplicial complexes with a cone*, *Comm. Algebra* **38** (2010), 3686–3698.
- [6] M. Barile and N. Terai, *The Stanley–Reisner ideals of polygons as set-theoretic complete intersections*, *Comm. Algebra* **39** (2011), 621–633.
- [7] R. Diestel, *Graph Theory*, 2nd edition, GTM **173** Springer, Berlin/Heidelberg/New York, 2000.
- [8] Brendan D. McKay, *NAUTY: No AUTomorphisms, Yes?*, Available at <http://cs.anu.edu.au/~bdm/nauty/>
- [9] CoCoATeam, *CoCoA: a system for doing Computations in Commutative Algebra*, Available at <http://cocoa.dima.unige.it>
- [10] T. Hibi, *Algebraic combinatorics on convex polytopes*, Carslaw Publications, Glebe, N.S.W., Australia, 1992.
- [11] K. Kimura, *Lyubeznik resolutions and the arithmetical rank of monomial ideals*, *Proc. Amer. Math. Soc* **137** (2009), 3627–3635.
- [12] K. Kimura, *Arithmetical rank of Cohen–Macaulay squarefree monomial ideals of height two*, *J. Commut. Algebra* **3** (2011), 31–46.
- [13] K. Kimura, G. Rinaldo and N. Terai, *Set of routines to generate and analyze all the squarefree monomial ideals with 5 generators*, Available at <http://ww2.unime.it/algebra/rinaldo/sqfreemonid/>, 2011.
- [14] K. Kimura, N. Terai and K. Yoshida, *Arithmetical rank of squarefree monomial ideals of small arithmetical degree*, *J. Algebraic Combin.* **29** (2009), 389–404.
- [15] K. Kimura, N. Terai and K. Yoshida, *Arithmetical rank of monomial ideals of deviation two*, in *Combinatorial Aspects of Commutative Algebra* (V. Ene and E. Miller eds.), *Contemporary Mathematics*, AMS, **502** (2009), 73–112.
- [16] M. Kummini, *Regularity, depth and arithmetic rank of bipartite edge ideals*, *J. Alg. Combin.* **30**(2009), 429–445.
- [17] G. Lyubeznik, *On the local cohomology modules $H_a^i(R)$ for ideals a generated by monomials in an R -sequence*, in *Complete Intersections*, Acireale, 1983 (S. Greco and R. Strano eds.), *Lecture Notes in Mathematics* No. 1092, Springer-Verlag, 1984, pp. 214–220.
- [18] G. Lyubeznik, *A new explicit finite free resolutions of ideals generated by monomials in an R -sequence*, *J. Pure Appl. Algebra* **51** (1988), 193–195.
- [19] M. Morales, *Simplicial ideals, 2-linear ideals and arithmetical rank*, *J. Algebra* **324** (2010), 3431–3456.
- [20] T. Schmitt and W. Vogel, *Note on set-theoretic intersections of subvarieties of projective space*, *Math. Ann.* **245** (1979), 247–253.
- [21] R. P. Stanley, *Linear diophantine equations and local cohomology*, *Inventiones mathematicae* **68**(1982) 175–193.
- [22] R.P. Stanley, *Combinatorics and commutative algebra*, 2nd edition, Birkäuser, Boston/Basel/ Stuttgart, 1996.
- [23] N. Terai, *Alexander duality in Stanley–Reisner rings*, in “Affine Algebraic Geometry, in honor of Professor Masayoshi Miyanishi on the Occasion of His Retirement from Osaka University (Hibi ed.)”, pp. 449–462, 2007, Osaka University Press.
- [24] Z. Yan, *An étale analog of the Goresky–Macpherson formula for subspace arrangements*, *J. Pure Appl. Algebra* **146** (2000), 305–318.

(Kyouko Kimura) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, SHIZUOKA UNIVERSITY, 836 OHYA, SURUGA-KU, SHIZUOKA 422-8529, JAPAN

E-mail address: `skkimur@ipc.shizuoka.ac.jp`

(Giancarlo Rinaldo) DIPARTIMENTO DI MATEMATICA, UNIVERSITA' DI MESSINA, SALITA SPERONE, 31. S. AGATA, MESSINA 98166, ITALY

E-mail address: `rinaldo@dipmat.unime.it`

(Naoki Terai) DEPARTMENT OF MATHEMATICS, FACULTY OF CULTURE AND EDUCATION, SAGA UNIVERSITY, SAGA 840-8502, JAPAN

E-mail address: `terai@cc.saga-u.ac.jp`